



Laparoscopic resection of choledochal cysts in adults: a series from Turkey

Ersin Gündoğan¹, Fatih Sümer¹, Muhammed Kadri Çolakoğlu², Egemen Çiçek¹, Cihan Gökler¹, Mehmet Can Aydın¹, Cüneyt Kayaalp¹

¹ Department of General Surgery, Inonu University School of Medicine, Malatya, Turkey

² Department of General Surgery, Recep Tayyip Erdoğan University School of Medicine, Rize, Turkey

ABSTRACT

Objective: Choledochal cyst is a congenital disease in which surgical treatment is preliminary because of the potential for malignancy. In recent years, increase in technological developments and laparoscopic experience have popularised the use of laparoscopy in adult choledochal cyst surgery. This study aimed to present the results of eight adult patients undergoing laparoscopic choledochal excision surgery.

Material and Methods: Patients who underwent laparoscopic choledochal cyst excision and hepatico-jejunostomy anastomoses between the years 2013 and 2018 were evaluated retrospectively. Demographic characteristics, preoperative and postoperative findings, pathological results and final condition of the patients were examined.

Results: Of the eight patients, three were males and five were females. Median age was 41.5 years (22-49). One of the patients had Type IVa and the rest had Type I choledochal cysts. Laparoscopic choledochal cyst excision, cholecystectomy, and hepatico-jejunostomy anastomoses were performed on all of the patients. One patient was converted to open surgery. Three patients had postoperative biliary leakage. Duration of the operations was determined as median 330 (240-480) minutes and blood loss was 50 (10-100) mL. Hospitalization of the patients was median 6 (4-23) days and follow-up time was median 20 (2-65) months. In the late period, cholangitis occurred in a patient who was treated with medical therapy and there was no mortality in the follow-up period.

Conclusion: We suggest that laparoscopic choledochal cyst excision in adults may be an alternative to open surgery due to the satisfactory results in the late period in spite of early problems like self-limiting bile leakage.

Keywords: Laparoscopy, Hepatico-jejunostomy, minimally invasive surgery, cyst excision

INTRODUCTION

Choledochal cyst is a congenital disease which presents with cystic dilatation of the biliary tract. The incidence is one in 100.000 and 80% presents in childhood (1,2). Although choledochal cysts were first classified by Alonso-Lej, original classification used today was categorised by Todani (3). According to this classification, there are five types of choledochal cysts and the most common one is type 1; followed by types 4, 5, 3 and 2 in decreasing frequency. Abdominal pain, jaundice and choledochal enlargement, which are usually identified as symptoms triad, are detected in only 6% of the patients (4). Treatment of choledochal cyst is surgical excision and bilioenteric reconstruction due to recurrent cholangitis, biliary stone formation, pancreatitis and risk of malignancy. Recently, minimally invasive approaches have been used in the treatment of choledochal cysts with acceptable morbidity and mortality (5).

Our aim in this study was to emphasize the availability of laparoscopic choledochal cyst excision.

MATERIAL and METHODS

Eight patients scheduled for laparoscopic cyst excision and bilio-enteric reconstruction due to choledochal cyst in our clinic between February 2013 and May 2018 were included into the study. The study was conducted in accordance with the Declaration of Helsinki. Patients were operated with their informed consent in technical details as previously described (6) (Figure 1 and 2). Patient demographics, body mass indexes, history of additional diseases, history of previous operations, types of cysts, operation

Cite this article as: Gündoğan E, Sümer F, Çolakoğlu MK, Çiçek E, Gökler C, Aydın MC, et al. Laparoscopic resection of choledochal cysts in adults: a series from Turkey. Turk J Surg 2019; 35 (3): 223-226.

Corresponding Author

Ersin Gündoğan

E-mail: ersingundogan@hotmail.com

Received: 12.10.2018

Accepted: 20.11.2018

Available Online Date: 23.09.2019

© Copyright 2019 by Turkish Surgical Society Available online at www.turkjsurg.com

DOI: 10.5578/turkjsurg.4331



Figure 1. Choledochal cyst.



Figure 2. Hepatico- jejunostomy.

times, amount of blood losses, length of hospital stay, early and late complications, pathology findings and final conditions of the cases were reviewed.

Statistical Analysis

Descriptive statistics were performed for all data. Values were reported in median and range or mean and percentages. The data was analyzed using Microsoft Excel 2013 program.

RESULTS

Sixty-two percent of the patients were females, and mean age was median 41.5 (22-49) years. There were no known comorbidities for all patients. There was a previous laparoscopic cholecystectomy history in one patient. One of the patients had Type IVa and the rest had Type I choledochal cysts. Choledochal excision and hepatico-jejunostomy anastomoses procedure were performed for all patients as surgical treatment. In all patients, surgical procedure was completed laparoscopically. However, after completing the surgical procedure, bile leak was detected from hepatico-jejunostomy anastomoses intraoperatively in a patient and converted to open surgical procedure for revision (14%). Hepatico-jejunostomy anastomosis of the patient was renewed and the operation ended after no problem was detected. Mean duration of the operation was median 330 (240-480) minutes and blood loss was median 50

(10-100) mL. All patients were followed for one day postoperatively in the intensive care unit, and no blood replacement was necessary for any patient. Postoperative bile leak developed in three patients. One of these patients required drainage with relaparoscopy and spontaneous cessation of the biliary leakage occurred in the follow-up. Drainage was discontinued on postoperative 3rd and 4th days in the other two patients with bile leak. Postoperative pneumonia developed in one patient, and the patient was discharged without any problems after medical treatment. Hospitalization time of our patients was median 6 (4-23) days. Examination of the pathologic lesions revealed that all patients were compatible with choledochal cyst while one patient had an additional gallbladder tumor. Liver resection and lymphadenectomy were performed in a second session to the patient whose tumor was T2. Pathologic examination revealed no tumoral tissue in the liver, and all of the lymph nodes were reactive. Follow-up period of the patients was median 20 (2-65) months. In follow-up, cholangitis occurred in one patient who was treated with medical therapy, and there was no mortality in the follow-up period (Table 1).

DISCUSSION

Although choledochal cysts are rare, it is a congenital disease in which surgical treatment has priority because of the risk for malignancy. It is usually seen in childhood, but 20% is detected in adult ages (2). It is detected more in females as in our study. In the surgical treatment of choledochal cysts, hepatico-jejunostomy or hepatico-duodenostomy anastomoses are preferred following choledochal excision. However, considering that pancreaticobiliary reflux after hepatico-duodenostomy anastomoses increases the risk of malignancy in the hepatic hilus, hepatico-jejunostomy anastomoses has become the preferred surgical procedure as it is in our study (7). Laparoscopic cyst excision and bilio-enteric reconstruction are used more frequently nowadays with the developments in minimally invasive surgery. With this application, it has been reported that abdominal wall complications (wound infection and incisional hernia) and perioperative blood loss are reduced. It has also been reported that laparoscopy provides a more detailed image due to its enlargement feature (8,9). However, the application of laparoscopy to biliary reconstructive surgery has been limited in part because of the complexity of these procedures and the need for rigorous surgical technique (8,10). To the best of our knowledge, this is the largest experience in minimally invasive surgery on choledochal cysts in Turkey.

Another limitation of the laparoscopic surgery practice is that it has longer operating times than open surgery. Our median operation time was 330 minutes, and it was reported in the literature with the range of 228 and 395 minutes (7-15). Liu et al. have reported that when compared with open surgery, the laparoscopic approach shows a longer operative time (249 ± 58 vs. 132 ± 15 min). Their study has revealed that time of laparos-

Table 1. Demographic, preoperative and postoperative findings

Age	Sex	History of abdominal operation	Type	Blood Loss (mL)	Operation Time (min)	Conversion	Complication	Hospital stay (day)	Follow up (month)	Morbidity
40	M	None	I	80	260	No	None	6	65.3	None
22	F	CIS	I	50	360	No	Pneumonia (2)	6	40.1	None
43	F	CIS Appendectomy	I	100	480	No	None	4	41.1	None
45	F	Cholecystectomy	I	70	240	Yes	None	8	23.8	None
23	F	None	I	50	380	No	Bile leak (2)	7	16.7	None
48	F	Appendectomy	I	10	420	No	None	4	12.8	None
49	M	None	IVa	30	300	No	Bile leak (3b)	23	10.2	Cholangitis
28	M	None	I	20	280	No	Bile leak (2)	6	1.8	None

Complication classification was made according to Clavien-Dindo Classification.

copric procedure decreases after twenty cases and lasts for up to 190 minutes (9). Therefore, they have shared the idea that after completing the learning curve, laparoscopic procedures would have similar operative times with open operations.

In our study, the amount of blood loss was median 50 mL. In the study by Liu et al., the laparoscopic group and open surgery group have been compared and the mean amount of blood loss was found as 72 ± 26 mL in the laparoscopic group and as 174 ± 51 mL in the open group (9). In a study by Senthilnathan et al., 110 patients undergoing laparoscopic surgery have been studied and mean blood loss was 66.2 ± 24.3 mL (8). Tian and colleagues have determined this amount as 252.3 ± 162.5 mL (13). Based on these results, it is seen that the amount of blood loss in our study is compatible with the literature.

In our study, length of hospitalization was median six days. In the study of Lu et al with 34 patients, it has been reported as 4.7 ± 1.3 days, and besides that, in the study of Jang and colleagues with 82 patients, it has been found as 8.6 ± 5.2 days and 8.3 ± 3.2 days in the study of Tian et al (11-13).

Although literature reveals bile leak rate between 3.6% and 12.8%, it is still the most common complication. However, it usually resolves without any intervention (8-15). The rate of bile leakage in our study seems to be higher than the literature (38.5%). However, two of the three cases with bile leak were spontaneously resolved on postoperative third and fourth days without any additional intervention.

Other complications mentioned in the literature include intra-abdominal collection, hemorrhage, pancreatic fistula, pancreatitis, intestinal obstruction, stricture in hepatico-jejunostomy anastomosis, and wound complications (12). In our series, only one patient (type IVa) was admitted to our hospital with late cholangitis and treated medically.

CONCLUSION

We suggest that laparoscopic choledochal cyst excision in adults may be an alternative to traditional open surgery due to the satisfactory results in the late period in spite of the early problems like self-limiting bile leakage.

Ethics Committee Approval: This study is registered at Inonu University Human Ethical Committee as a research. The registration identification number is 2014/33.

Informed Consent: Written informed consent was obtained from patients who participated in this study.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - E.G.; Design - E.G., F.S.; Supervision - F.S., C.K.; Resource - E.G., M.K.Ç.; Materials - E.G., M.K.Ç.; Data Collection and/or Processing - E.Ç.; Analysis and/or Interpretation - C.G.; Literature Search - M.C.A.; Writing Manuscript - E.G., F.S.; Critical Reviews - C.K.

Conflict of Interest: The authors have no conflicts of interest to declare.

Financial Disclosure: The authors have no conflicts of interest to declare.

REFERENCES

- Gigot J, Nagorney D, Farnell M, Moir C, Ilstrup D. Bile duct cysts: a changing spectrum of disease. *J Hepatobiliary Pancreat Surg* 1996;3:405-11. [\[CrossRef\]](#)
- Stain SC, Guthrie CR, Yellin AE, Donovan AJ. Choledochal cyst in the adult. *Ann Surg* 1995;222:128-33. [\[CrossRef\]](#)
- Dumitrascu T, Lupescu I, Ionescu M. The Todani classification for bile duct cysts: an overview. *Acta Chir Belg* 2012;112:340-5. [\[CrossRef\]](#)
- Stringer MD, Dhawan A, Davenport M, Mieli-Vergani G, Mowat AP, Howard ER. Choledochal cysts: lessons from a 20 year experience. *Arch Dis Child* 1995;73:528-31. [\[CrossRef\]](#)
- Lipsett PA, Pitt HA, Colombani PM, Boitnott JK, Cameron JL. Choledochal cyst disease: a changing pattern of presentation. *Ann Surg* 1994;220:644-52. [\[CrossRef\]](#)
- Kayaalp C, Soyer V, Ersan V, Aydın C, Karagul S. Laparoscopic choledochal cyst excision and Roux-en-Y hepaticojejunostomy. *Turk J Surg* 2015;32:152-4. [\[CrossRef\]](#)

7. Singham J, Yoshida EM, Sendamore CH. Choledochal cysts. Part 3 of 3: management. *Can J Surg* 2010;53:51-6.
8. Senthilnathan P, Patel ND, Nair AS, Nalankilli VP, Vijay A, Palanivelu C. Laparoscopic management of choledochal cyst-technical modifications and outcome analysis. *World J Surg* 2015;39:2550-6. [CrossRef]
9. Liu Y, Yao X, Li S, Liu W, Liu L, Liu J. Comparison of therapeutic effects of laparoscopic and open operation for congenital choledochal cysts in adults. *Gastroenterol Res Pract* 2014;2014:1-8. [CrossRef]
10. Palanivelu C, Rangarajan M, Parthasarathi R, Amar V, Senthilnathan P. Laparoscopic management of choledochal cysts: technique and outcomes: a retrospective study of 35 patients from a tertiary center. *J Am Coll Surg* 2008;207:839-46. [CrossRef]
11. Lü SC, Shi XJ, Wang HG, Lu F, Liang YR, Luo Y, et al. Technical points of total laparoscopic choledochal cyst excision. *Chin Med J (Engl)* 2013;126:884-7.
12. Jang JY, Yoon YS, Kang MJ, Kwon W, Park JW, Chang YR, et al. Laparoscopic excision of a choledochal cyst in 82 consecutive patients. *Surg Endosc Other Interv Tech* 2013;27:1648-52. [CrossRef]
13. Tian Y, Wu SD, Zhu AD, Chen DX. Management of type I choledochal cyst in adult: totally laparoscopic resection and Roux-en-Y hepaticoenterostomy. *J Gastrointest Surg* 2010;14:1381-8. [CrossRef]
14. Jang JY, Kim SW, Han HS, Yoon YS, Han SS, Park YH. Totally laparoscopic management of choledochal cysts using a four-hole method. *Surg Endosc Other Interv Tech* 2006;20:1762-5. [CrossRef]
15. Hwang DW, Lee JH, Lee SY, Song DK, Hwang JW, Park KM, et al. Early experience of laparoscopic complete en bloc excision for choledochal cysts in adults. *Surg Endosc Other Interv Tech* 2012;26:3324-9. [CrossRef]



ORJİNAL ÇALIŞMA-ÖZET
Türk J Surg 2019; 35 (3): 223-226

Erişkinlerde laparoskopik koledok kist rezeksiyonu: Türkiye'den bir seri

Ersin Gündoğan¹, Fatih Sümer¹, Muhammed Kadri Çolakoğlu², Egemen Çiçek¹, Cihan Gökler¹, Mehmet Can Aydın¹, Cüneyt Kayaalp¹

¹ İnönü Üniversitesi Tıp Fakültesi, Genel Cerrahi Anabilim Dalı, Malatya, Türkiye

² Recep Tayyip Erdoğan Üniversitesi Tıp Fakültesi, Genel Cerrahi Anabilim Dalı, Rize, Türkiye

ÖZET

Giriş ve Amaç: Koledok kistleri malignite potansiyeli nedeniyle cerrahi tedavinin ön planda olduğu konjenital bir hastalıktır. Son yıllarda teknolojik gelişmeler ve laparoskopik tecrübeye artış erişkin koledok kist cerrahisinde laparoskopinin kullanımını yaygınlaştırmaktadır. Bu çalışmada amacımız laparoskopik koledok eksizyonu yapılan sekiz erişkin hastanın sonuçlarını sunmaktır.

Gereç ve Yöntem: Kliniğimizde 2013-2018 yılları arasında laparoskopik koledok kist eksizyonu ve hepatiko-jejunostomi uygulanan hastalar retrospektif olarak değerlendirildi. Hastaların demografik özellikleri, peroperatif ve postoperatif bulguları, patolojik sonuçları ve son durumları irdelendi.

Bulgular: Çalışmaya dahil edilen sekiz hastanın üçü erkek, beşi kadın idi. Yaşların median değeri 41.5 (22-49) idi. Hastaların bir tanesinde tip IVa, diğerlerinde ise tip 1 koledok kisti saptandı. Hastaların tamamına laparoskopik koledok eksizyonu, kolesistektomi ve hepatiko-jejunostomi yapıldı. Bir hastada açık cerrahiye geçildi. Üç hastada postoperatif safra kaçağı oldu. Operasyon süreleri median 330 (240-480) dakika ve kan kaybı 50 (10-100) mL olarak saptandı. Hastaların yatış süreleri median 6 (4-23) gün ve takip süreleri median 20 (2-65) ay olduğu tespit edildi. Geç dönemde; bir hastada medikal tedavi ile düzelen kolanjit atağı gözlemlendi, mortalite ise görülmedi.

Sonuç: Kendini sınırlayan safra kaçaqları dışında erken dönemde önemli sorun yaşamamamız ve geç dönemdeki tatminkar sonuçlar nedeniyle laparoskopik koledok kist eksizyonunun erişkinlerde açık cerrahiye bir alternatif olabileceğini söyleyebiliriz.

Anahtar Kelimeler: Laparoskopi, hepatiko-jejunostomi, minimal invaziv cerrahi, koledok kist eksizyonu

DOI: 10.5578/turksurg.4331