

1-1-2010

## A new study on the bacteria of *Agrotis segetum* Schiff. (Lepidoptera: Noctuidae) and their insecticidal activities

ALİ SEVİM

ZİHNİ DEMİRBAĞ

İSMAİL DEMİR

Follow this and additional works at: <https://journals.tubitak.gov.tr/agriculture>



Part of the [Agriculture Commons](#), and the [Forest Sciences Commons](#)

---

### Recommended Citation

SEVİM, ALİ; DEMİRBAĞ, ZİHNİ; and DEMİR, İSMAİL (2010) "A new study on the bacteria of *Agrotis segetum* Schiff. (Lepidoptera: Noctuidae) and their insecticidal activities," *Turkish Journal of Agriculture and Forestry*. Vol. 34: No. 4, Article 7. <https://doi.org/10.3906/tar-0902-9>  
Available at: <https://journals.tubitak.gov.tr/agriculture/vol34/iss4/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Agriculture and Forestry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## A new study on the bacteria of *Agrotis segetum* Schiff. (Lepidoptera: Noctuidae) and their insecticidal activities

Ali SEVİM<sup>1</sup>, Zihni DEMİRBAĞ<sup>2</sup>, İsmail DEMİR<sup>2,\*</sup>

<sup>1</sup>Department of Biology, Faculty of Arts and Sciences, Rize University, 53100 Rize - TURKEY

<sup>2</sup>Department of Biology, Faculty of Arts and Sciences, Karadeniz Technical University, 61080 Trabzon - TURKEY

Received: 11.02.2009

**Abstract:** *Agrotis segetum* Schiff. (Lepidoptera: Noctuidae) is one of the most serious pests of nearly all vegetables in Turkey. In this study, to find a more effective and safe biological control agent, the bacterial flora of *Agrotis segetum* were investigated in larvae collected from different populations in the vegetable fields of the Black Sea region of Turkey, and 9 bacterial isolates were cultured from these populations. Seven of these bacteria were determined and characterized at the species level and the rest were characterized at the genus level. Morphological, physiological, and biochemical characteristics of the bacterial isolates were determined by conventional and routine techniques. Characterization of the isolates was also performed using the VITEK 32 bacterial identification system. In addition, a 16S rRNA gene sequence was determined for these isolates to query against the NCBI genetic database and to construct a phylogenetic tree using closely related species. Consequently, the isolates were identified as *Bacillus cereus* (Ags1), *Bacillus* sp. (Ags2), *Bacillus megaterium* (Ags3), *Enterobacter aerogenes* (Ags4), *Acinetobacter calcoaceticus* (Ags5), *Enterobacter* sp. (Ags6), *Pseudomonas putida* (Ags7), *Enterococcus gallinarum* (Ags8), and *Stenotrophomonas maltophilia* (Ags9). All isolates were determined the first time from the flora of *A. segetum*. Furthermore, the insecticidal activities of the bacterial isolates were tested against *A. segetum* larvae, and *Enterococcus gallinarum* (Ags8) was found to cause the highest insecticidal activity (60%) 8 days after application.

**Key words:** *Agrotis segetum*, bacterial flora, insecticidal activity

### *Agrotis segetum* Schiff. (Lepidoptera: Noctuidae)'un bakterileri ve bu bakterilerin insektisidal etkileri üzerine yeni bir çalışma

**Özet:** *Agrotis segetum* Schiff. (Lepidoptera: Noctuidae) Türkiye'de hemen hemen bütün sebzelerin en önemli zararlılarından birisidir. Bu çalışmada, daha etkili ve güvenli biyolojik mücadele etmeni bulmak için *Agrotis segetum*'un bakteriyel florası Karadeniz Bölgesi'ndeki sebze tarlalarında farklı popülasyonlardan toplanan larvalar üzerinde araştırıldı. Bu popülasyonlardan dokuz bakteri izolatu kültüre alındı. Bu bakterilerden yedi tanesi tür seviyesinde, diğerleri ise cins seviyesinde tanımlandı ve karakterize edildi. Bakteriyel isolatların morfolojik, fizyolojik ve biyokimyasal özellikleri, rutin ve geleneksel teknikler kullanılarak belirlendi. İzolatların karakterizasyonu, VITEK 32 bakteriyel tanımlama sistemi kullanılarak da gerçekleştirildi. Buna ilaveten, bütün isolatların NCBI genetik veritabanında karşılaştırmalarını yapmak ve ilişkili türler ile filogenetik yakınlıklarını belirlemek için 16S rRNA gen sekans analizleri yapıldı. Sonuç olarak, isolatlar *Bacillus cereus* (Ags1), *Bacillus* sp. (Ags2), *B. megaterium* (Ags3), *Enterobacter aerogenes* (Ags4), *Acinetobacter calcoaceticus* (Ags5), *Enterobacter* sp. (Ags6), *Pseudomonas putida* (Ags7), *Enterococcus gallinarum* (Ags8) ve *Stenotrophomonas maltophilia* (Ags9) olarak tanımlandı. Bütün isolatlar *A. segetum*'un florasında ilk kez belirlendi. Ayrıca, bakteriyel isolatların insektisidal etkileri *A. segetum* larvalarına karşı test edildi. Uygulamadan 8 gün sonra *Enterococcus gallinarum* (Ags8)'un en yüksek aktiviteye (%60) sebep olduğu bulundu.

**Anahtar sözcükler:** *Agrotis segetum*, bakteriyel flora, insektisidal etki

\* E-mail: idemir@ktu.edu.tr

## Introduction

Cutworms are important pests for a number of crops in agriculture, horticulture, and forestry around the world (Zethner 1980). The common cutworm (turnip moth), *Agrotis segetum* Schiff. (Lepidoptera: Noctuidae), is a serious pest occurring throughout Europe (including Turkey), Asia, and parts of Africa (Jakubowska et al. 2005). It commonly lives on the ground, where it feeds on seedlings of nearly all vegetable and field crops, most importantly corn, potatoes, beans, peppers, eggplants, okra, lettuce, tobacco, sugar beets, and cabbage, as well as many other plants (The Ministry of Agriculture of Turkey 2008).

Chemical substances such as chlorpyrifos and endosulfan have been utilized to control this pest in Turkey (The Ministry of Agriculture of Turkey 2008). However, recent concerns about the hazardous effects of chemical pesticides on the environment have encouraged scientists to consider finding more effective and safe control agents. Use of some microbial control agents against *A. segetum* was therefore previously studied. Microbial insecticides based on entomopathogenic bacteria, viruses, nematodes, and fungi have been assessed as possible microbial control agents of *A. segetum* (Lipa and Wiland 1972; Ignoffo and Garcia 1979; Lossbroek and Theunissen 1985; Caballero et al. 1991; Steenberg and Ogaard 2000; Yun et al. 2004).

Studies of symbiotic bacteria in insect species allow the development of new approaches for biological control. Symbiotic bacteria are ubiquitously located in animal guts with these symbioses ranging from pathogenic to mutualistic and from facultative to obligate (Lau et al. 2002). In the last few years, several pathogenic bacteria species from insect samples have been developed as pesticides and used successfully in biological control of insects worldwide (Thiery and Frachon 1997). Several studies have been performed to determine the bacterial flora of *A. segetum* (Lipa and Wiland 1972; Turchaninova 1972; Turchaninova 1973), but no such study has been done in Turkey.

Efforts have been increased to discover more effective and safe biological control against hazardous insects to replace potentially hazardous chemical agents (Ince et al. 2008). Moreover, investigating a new pathogen from various environments against a

target pest is always desirable. It is also known that the success of control can be higher when native isolates are used against the target insect. Therefore, to find a more effective and safe biocontrol agent against cutworm, we investigated the bacterial flora of *A. segetum* isolated from the Black Sea region of Turkey, utilizing current morphological, biochemical, and molecular techniques. In this study, we isolated 9 bacterial isolates, characterized them in detail, and tested their insecticidal activities against larvae of *A. segetum*.

## Materials and methods

### Collection of larvae

Larvae of the cutworm, different instars, were used for the isolation of bacteria. Larvae were collected on a daily basis from different tobacco fields in the vicinity of Samsun in May and June of 2007. Larvae were obtained by digging them out of the soil, and they were placed individually into plastic boxes (20 mm) with a little soil and perforated covers to permit airflow. The boxes were transported to the laboratory. Larvae were fed with lettuce at room temperature with a 12:12 photoperiod until the bacterial isolation was done. The lettuce was washed with sterile water and dried before being given to the larvae.

The larvae used in the insecticidal activity tests were provided by Dr. Christer Löfstedt and Erling Jirle (Department of Ecology, Lund University, Sweden) as egg mass.

### Isolation and purification of bacterial samples

After macroscopic examination, living larvae that showed general disease symptoms were separated from dead larvae. After that, both dead and living larvae were surface sterilized with 70% alcohol for 2 min and then washed twice with sterile distilled water. The larvae were homogenized in nutrient broth using a glass tissue grinder. The obtained larval extracts were used to prepare a series of dilutions, from  $10^{-1}$  to  $10^{-5}$ . After preparation of the dilutions, 2 methods were used for bacterial purification. In the first, the larval suspension was heated at 80 °C for 10 min in a water bath to eliminate non-spore-forming organisms (Ohba and Aizawa 1986). After that, 100 µL of this suspension was streaked on each nutrient agar (Difco) plate and incubated at 30 °C for 2-3 days. In the

second method, serially diluted suspensions were also spread on nutrient agar (Difco) plates and incubated under the same conditions. At the end of the incubation period, isolates were separated based on the color and morphology of the colonies. Pure cultures of bacterial colonies were then prepared, and these cultures were identified by their morphology, spore formation, and nutritional features; their physiological, biochemical, and molecular characteristics; and by the VITEK 32 bacterial identification system.

#### Identification of bacterial isolates based on conventional tests

Morphological, physiological, and biochemical properties of the bacterial isolates were determined according to *Bergey's Manual of Systematic Bacteriology*, Volumes 1 and 2 (Palleroni 1984; Kandler and Weiss 1986). Gram staining was performed according to the procedure described by Claus (1992). Endospore staining was performed according to the method of Prescott et al. (1996). Capsule staining was performed by negative staining. The motility of isolates was determined by using a semisolid medium (Soutourina et al. 2001). Physiological tests were performed by using different media and the VITEK 32 bacterial identification system. The pH and NaCl tolerance values were determined in LB broth.

#### Isolate identification by VITEK 32

The stock culture strains were then subcultured onto tryptic soy agar plates to check their purity. A standard bacterial suspension was prepared in 1.8 mL of 0.45% saline using the VITEK colorimeter for each isolate. The time interval between suspension preparation and card filling was less than 30 min to avoid changes in turbidity. After preparation of the suspension, it was inoculated into the VITEK GNI (for gram-negative bacteria), GPI (for gram-positive bacteria), and CAP (for rod-shaped bacillus bacteria) cards, incubated at 35 °C in the reader incubator module for 18 h, and automatically read hourly by the optical scanner according to the manufacturer recommendations by using bioLiaison software. In addition, since the VITEK system requires the oxidase (gram-negative) and catalase (gram-positive) test results for identification, we also determined features of oxidase and catalase tests (Barry et al. 1982).

#### Isolate identification by 16S rRNA gene sequencing

Isolation of genomic DNA was performed according to standard phenol/chloroform procedures (Sambrook et al. 1989). DNA pellets were dissolved in 50 µL of TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0) and isolated DNAs were stored at -20 °C until use. PCR amplification of the 16S rRNA gene of bacterial isolates was performed with the following primers: Forward UNI16S-L, 5'-ATTCTAGAGTTTGATCATGGCTCA-3'; and Reverse UNI16S-R, 5'-ATGGTACCGTGTGACGGGCGGTGTGTA-3' (William et al. 1991). PCR conditions were adapted essentially as described by William et al. (1991). Gel-purified PCR products of the 16S rRNA gene fragments from the 9 isolates were cloned directly into the pGEM-T vector cloning system and transformed into *E. coli* JM 101 strain. After that, plasmid isolation was done and plasmid DNA samples were sent to Macrogen for sequencing. The obtained sequences data were submitted to GenBank for accession numbers and were compared with known 16S rRNA gene sequences in the GenBank database using the BLAST search algorithm. Finally, the taxonomic identification of bacterial isolates was determined by comparing the results obtained by conventional tests, VITEK 32, and 16S rRNA gene analysis data.

#### Insecticidal effects of bacterial isolates

Experimental infections were carried out for the isolates to determine their insecticidal activities. *A. segetum* eggs were obtained from laboratory cultures from Sweden (Department of Ecology, Lund University) and brought to the laboratory. Hatched larvae were grown in the laboratory as described above. Healthy larvae were used for the insecticidal effects of bacterial isolates. Isolates were incubated for 18 h (72 h for *Bacillus* for sporulation) at 30 °C in nutrient broth. After incubation, the density of cells was set at 1.89 at OD<sub>600</sub>, and 5 mL of culture was centrifuged at 3000 rpm for 10 min (Ben-Dov et al. 1995). The pellet was resuspended in 5 mL of sterilized PBS and used for bioassays. Fresh leaves of lettuce used as food were dipped into the bacterial suspension and placed into individual sterile plastic containers (80 mm in diameter), each containing a single bacterial isolate. For the control group, leaves of lettuces were dipped into sterile water. Ten third instars larvae were placed on the lettuce in containers and kept at 26 ± 2 °C and 60% relative humidity with a 12:12 photoperiod (Mitchell and Smith 1985). The mortalities of larvae were

recorded every 24 h, with all dead larvae removed from the containers. Data were evaluated by using Abbott's formula (Abbott 1925).

**Phylogenetic analysis of bacterial isolates**

The evolutionary relationship of the 9 bacterial isolates and 28 closely related species were evaluated. An approximately 1400 bp segment of 16S rRNA was used for phylogenetic analysis. Sequences were assembled and edited with BioEdit and aligned (Hall 1999). Cluster analyses of the sequences were performed using BioEdit (version 7.09) with ClustalW. To evaluate the phylogenetic relationship of isolates, neighbor-joining analysis was performed on aligned sequences by using the Kimura 2-parameter test with MEGA 4.0 software (Tamura et al. 2007). Alignment gaps were treated as missing data. The reliability of the phylograms was tested by bootstrap analysis with 1000 replicates using MEGA 4.0.

**Results**

A total of 9 isolates were finally selected and characterized according to morphology, spore formation, nutritional features, physiological and biochemical characteristics, results of the VITEK 32 system, and analysis of the 16S rRNA gene sequence (Tables 1, 2, 3, 4, and 5).

On agar plates, 8 isolates were cream-colored and 1 isolate was orange (Table 1). Colonies of all isolates were smooth, except for Ags1, which was filamentous after 1 day of incubation on a nutrient agar plate at 30 °C. Two spore-forming bacteria (Ags1 and Ags3) and 7 non-spore-forming bacteria were identified. Seven isolates were rod-shaped and 2 isolates were coccobacillus. Four isolates (Ags1, Ags2, Ags3, and Ags8) were gram-positive and the rest of the isolates were gram-negative. Motility was observed in 6 isolates (Ags1, Ags4, Ags5, Ags6, Ags7, and Ags8). All

Table 1. The morphological characteristics of the bacterial isolates.

Isolate number	Ags1	Ags2	Ags3	Ags4	Ags5	Ags6	Ags7	Ags8	Ags9
Color of colonies	Cream	Orange	Cream	Cream	Cream	Cream	Cream	Cream	Cream
Shape of colonies	Irregular	Smooth	Smooth	Smooth	Smooth	Smooth	Round	Smooth	Smooth
Gram stain	+	+	+	-	-	-	-	+	-
Shape of bacteria	Bacillus	Bacillus	Bacillus	Bacillus	Coccobacillus	Coccobacillus	Bacillus	Bacillus	Bacillus
Spore stain	+	-	+	-	-	-	-	-	-
Spore shape	ND <sup>b</sup>	ND	ND	ND	ND	ND	ND	ND	ND
Spore form	Central	-	Central	-	-	-	-	-	-
Capsule	+	+	+	+	+	+	+	+	+
Turbidity when grown in NB <sup>a</sup>	Turbid	Turbid	Turbid	Turbid	Turbid	Turbid	Turbid	Turbid	Turbid
Motility	+	-	-	+	+	+	+	+	-

<sup>a</sup>NB: Nutrient Broth; <sup>b</sup>ND: No Data

Table 2. The physiological characteristics of the bacterial isolates.

Isolate number	Ags1	Ags2	Ags3	Ags4	Ags5	Ags6	Ags7	Ags8	Ags9
Growth in 2% NaCl	+	+	+	+	+	+	+	-	+
Growth in 5% NaCl	-	-	+	+	-	+	+	-	-
Growth in 7% NaCl	-	-	+	+	-	+	-	-	-
Growth in 12% NaCl	-	-	-	-	-	-	-	-	-
Growth in pH 3	-	-	-	-	-	-	-	-	-
Growth in pH 4	-	-	-	+	+	+	+	-	-
Growth in pH 5	+	-	+	+	+	+	+	+	+
Growth in pH 5.5	+	-	+	+	+	+	+	+	+
Growth in pH 6	+	+	+	+	+	+	+	+	+
Growth in pH 7	+	+	+	+	+	+	+	+	+
Growth in pH 8	+	+	+	+	+	+	+	+	+
Growth in pH 9	+	+	+	+	+	+	+	+	+
Growth in pH 10	-	-	-	+	-	-	-	+	+
Growth in pH 12	-	-	-	-	-	-	-	-	-

Table 3. The biochemical characteristics of the bacterial isolates.

Isolate number	Ags1	Ags2	Ags3	Ags4	Ags5	Ags6	Ags7	Ags8	Ags9
Nitrate reduction	+	+	+	+	+	+	+	+	+
KIA	Basic	Basic	Basic	Basic	Basic	Basic	Basic	Acid	Basic
Oxidase	+	-	-	+	+	-	+	-	+
Methyl red test	+	ND <sup>a</sup>	+	-	+	+	-	+	-
Voges Proskauer	-	ND	-	-	-	-	-	-	-
Citrate	-	-	-	+	+	-	+	-	-
Hydrolysis of urea	+	+	+	+	+	+	+	+	+
Indol	-	-	-	-	-	+	-	ND	-
Gelatin hydrolysis	+	-	-	-	-	-	-	-	+
Catalase	+	+	+	+	+	+	+	-	+
Coagulase			-						
Hydrolysis of starch	+	+	+	+	+	-	-	-	-
Sucrose	+	-	+	+	-	-	-	+	-
Esculin	+	-	+	+	-	+	-	+	+
Lactose (10%)	ND	-	ND	+	+	+	-	+	+
Trehalose	+	ND	+	ND	ND	ND	ND	+	ND
Arginine	ND	-	ND	-	-	-	+	+	-
Mannitol	-	-	+	+	-	+	-	+	-
Arabinose	-	-	+	+	+	+	+	+	-
Urea	ND	-	ND	-	-	-	-	-	-
Raffinose	-	-	+	+	-	-	-	+	-
Ribose	-	ND	-	ND	ND	ND	ND	-	ND
Xylose	-	-	-	-	+	-	+	+	+
Sorbitol	-	-	+	+	-	-	-	-	-
DP 300	ND	-	ND	-	-	-	+	ND	-
Maltose	+	-	+	+	-	+	-	ND	+
Inositol	-	-	-	-	-	-	-	ND	-
TLA (10% lactose)	ND	-	ND	+	+		-	ND	-
Glucose (oxidative)	ND	-	ND	+	+	+	+	ND	-
Adonitol	ND	-	ND	+	-	-	-	ND	-
Glucose	+	-	+	+	-	+	-	ND	-
Growth control	ND	+	ND	+	+	+	+	ND	-
Malonate	ND	-	ND	+	+	+	+	ND	+
Coumaric acid	ND	-	ND	-	+	-	+	ND	-
Acetamide	ND	-	ND	-	-	-	+	ND	-
Plant indican	ND	-	ND	-	-	+	-	ND	+
Rhamnose	ND	-	ND	+	-	+	-	ND	-
The mark of oxidase	ND	-	ND	+	+	-	+	ND	+

<sup>a</sup>ND: No Data

Note: The 1st through 28th tests were conventional; the rest were performed with VITEK 32.



Table 4. Results of identification of isolates according to VITEK 32 and 16S rRNA.

Isolate Number	VITEK 32 System	16S Gene Sequence	16S Gene Similarity (%)	Accession Numbers
Ags1	91% <i>Bacillus cereus</i>	<i>Bacillus thuringiensis</i> serovar <i>kurstaki</i> BnBt	99%	EU153549
		<i>Bacillus weihenstephanensis</i> 32	99%	AB334765
		<i>Bacillus cereus</i> strain AH 521	99%	AF290554
		<i>Bacillus mycoides</i> CIP 103472	99%	AM747229
		<i>Bacillus thuringiensis</i> 2000031482	99%	AY138290
Ags2	99% <i>Acinetobacter lwoffii/junii</i>	<i>Bacillus arsenicus</i> strain S8-06	99%	EU620414
		<i>Bacillus</i> sp. JL872	99%	DQ985053
		<i>Bacillus barbaricus</i> VII-B3-A2	98%	AJ422145
		<i>Bacillus gelatini</i> 506	94%	DQ350818
Ags3	99% <i>Bacillus megaterium</i>	<i>Bacillus megaterium</i> P2P5	99%	EU221388
		<i>Bacillus subtilis</i> C14-1	99%	EU257452
		<i>Bacillus flexus</i> L2S2	99%	EU221413
Ags4	Unidentified Organism	<i>Enterobacter asburiae</i> J2S4	99%	EU221358
		<i>Enterobacter aerogenes</i> WAB1906	98%	AM184247
		<i>Enterobacter</i> sp. 3-1t	99%	EU543690
		<i>Pantoea agglomerans</i> WAB1872	99%	AM184214
		<i>Citrobacter freundii</i> strain GM1	99%	DQ133536
		<i>Enterobacter cloacae</i> 766	98%	AM778415
		<i>Erwinia aphidicola</i> GTC 1688	99%	AB273744
Ags5	99% <i>Acinetobacter calcoaceticus-baumannii</i>	<i>Acinetobacter calcoaceticus</i> HPC253	99%	AY346313
		<i>Acinetobacter rhizosphaerae</i> PhyCEm-460	99%	AM921638
		<i>Acinetobacter baumannii</i> Ab8	99%	AY847284
Ags6	34% <i>Enterobacter cancerogenus</i>	<i>Pantoea agglomerans</i> HDDMN03	99%	EU879089
		<i>Enterobacter</i> sp. TUT1014	98%	AB098582
		<i>Citrobacter</i> sp. BinzhouCLT	98%	AF530068
		<i>Enterobacter aerogenes</i> zjs04	98%	DQ857896
Ags7	96% <i>Pseudomonas fluorescens/putida</i>	<i>Pseudomonas plecoglossicida</i> FPC951	99%	AB009457
		<i>Pseudomonas putida</i> ATCC 17390	99%	AF094737
		<i>Pseudomonas taiwanensis</i> BCRC 17751	99%	EU103629
Ags8	91% <i>Enterococcus gallinarum</i>	<i>Enterococcus gallinarum</i> LMG 13129	99%	AJ301833
		<i>Enterococcus casseliflavus</i> strain F32	99%	EU151766
		<i>Enterococcus flavescens</i> CECT4481T	99%	AJ420802
		<i>Enterococcus saccharolyticus</i>	99%	ESU30931
Ags9	70% <i>Stenotrophomonas maltophilia</i>	<i>Xanthomonas retroflexus</i> BBTR24	99%	DQ337602
		<i>Stenotrophomonas maltophilia</i> S-1	99%	AB194708

Table 5. Taxonomic identification of bacterial isolates of *A. segetum*.

Isolate Number	Suggested identification	Accession numbers of 16S gene
Ags1	<i>Bacillus cereus</i>	FJ380120
Ags2	<i>Bacillus</i> sp.	FJ380121
Ags3	<i>Bacillus megaterium</i>	FJ380122
Ags4	<i>Enterobacter aerogenes</i>	FJ380123
Ags5	<i>Acinetobacter calcoaceticus</i>	FJ380124
Ags6	<i>Enterobacter</i> sp.	FJ380125
Ags7	<i>Pseudomonas putida</i>	FJ380126
Ags8	<i>Enterococcus gallinarum</i>	FJ380127
Ags9	<i>Stenotrophomonas maltophilia</i>	FJ380128

isolates caused turbidity in the nutrient broth medium.

The physiological and biochemical characteristics of the isolates are reported in Tables 2 and 3.

For a more detailed identification of these bacterial isolates, the VITEK 32 identification system was used and a 16S rRNA gene sequence analysis was performed (Table 4). According to the VITEK 32 identification system, among the 9 isolates tested, 8 isolates were correctly identified at the species level. Only one isolate could not be identified.

After analyzing the conclusions of the conventional tests, the VITEK 32 identification system, and the 16S rRNA gene sequences, we were able to identify 7 isolates at the species level [Ags1 (*Bacillus cereus*), Ags3 (*Bacillus megaterium*), Ags4 (*Enterobacter aerogenes*), Ags5 (*Acinetobacter calcoaceticus*), Ags7 (*Pseudomonas putida*), Ags8 (*Enterococcus gallinarum*), and Ags9 (*Stenotrophomonas maltophilia*)] and were able to identify 2 isolates at the genus level [Ags2 (*Bacillus* sp.) and Ags6 (*Enterobacter* sp.)] (Table 5). Figure 1 shows the phylogenetic analysis of the bacterial isolates and closely related species based on the 16S rRNA gene sequencing.

In addition, we tested the insecticidal activity of the bacterial isolates against third instar *A. segetum* larvae. Isolate Ags8 showed the highest insecticidal activity with 60% mortality 8 days after application (Figure 2). Isolates Ags1, Ags2, Ags3, Ags4, Ags5,

Ags6, Ags7, and Ags9 had 30%, 10%, 40%, 10%, 20%, 30%, 20%, and 30% mortality 8 days after application, respectively (Figure 2).

## Discussion

In this study, we characterized 9 bacterial isolates based on morphological, nutritional, and physiological characteristics; biochemical studies; and the VITEK 32 system and 16S rRNA sequences. Based on these tests, the bacterial flora of *Agrotis segetum* consisted of *Bacillus cereus* (Ags1), *Bacillus* sp. (Ags2), *B. megaterium* (Ags3), *Enterobacter aerogenes* (Ags4), *Acinetobacter calcoaceticus* (Ags5), *Enterobacter* sp. (Ags6), *Pseudomonas putida* (Ags7), *Enterococcus gallinarum* (Ags8), and *Stenotrophomonas maltophilia* (Ags9) (Table 5). For bacterial identification, conventional results are sufficient for identifying a new bacterial isolate most of the time, but occasionally it is very difficult to identify some bacteria based only on conventional tests (Bahar and Demirbag 2007). Hence, we supported identifications of bacterial isolates by using the VITEK 32 bacterial identification system and 16S rRNA gene sequence analysis. Some strains require further study for detailed characterization.

Although Lipa and Wiland (1972) determined 8 bacterial isolates from *A. segetum* larvae, including *Aerobacter cloacae*, *Aerobacter* sp., *Bacillus thrungiensis*, *Brevibacterium maris*, *Escherichia coli*, *Escherichia freundii*, *Micrococcus luteus*, and *Pseudomonas fluorescens*, all of the isolates in our study were determined for the first time from the bacterial flora of *A. segetum*. Three species were determined to belong to the genus *Bacillus*. These isolates were identified as *Bacillus cereus* (Ags1), *Bacillus* sp. (Ags2), and *B. megaterium* (Ags3). The levels of 16S rRNA sequence similarity among the isolates are shown in Table 4. The differences between isolates of previous studies and this study are probably due to insufficient identification methods utilized in previous studies. Unfortunately, it is not possible to compare these isolates because there is no culture collection from the previous studies.

A common soil organism, *Bacillus cereus*, has been found to be a pathogen for insects on several occasions, and this species has been isolated from






Figure 1. Phylogenetic analysis of bacterial isolates from *A. segetum* and closely related species based on 16S rRNA gene sequencing. Bootstrap values are shown next to nodes and are based upon 1000 replicates using MEGA 4.0.

several insect species (Kuzina et al. 2001; Sezen et al. 2005). Like *B. cereus*, *B. megaterium* is common in insect populations (Osborn et al. 2002). So far, several *Bacillus* species have been isolated from different insect species and are very common in insect populations (Demir et al. 2002; Murrell et al. 2003; Yilmaz et al. 2006; Ince et al. 2008).

Like the genus *Bacillus*, several species of *Enterobacter* have been isolated from several insect

species and used for biological control purposes (Sandra and Douglas 2004; Bahar and Demirbag 2007). Other isolates determined in this study [*Acinetobacter calcoaceticus* (Ags5), *Pseudomonas putida* (Ags7), *Enterococcus gallinarum* (Ags8), and *Stenotrophomonas maltophilia* (Ags9)] have also already been shown to be common in insect populations (Yilmaz et al. 2006; Bahar and Demirbag 2007; Cox and Gilmore 2007).




Figure 2. Mortality of the third instar larvae of *Agrotis segetum* 8 days after application of bacterial isolates. Mortality data were corrected according to Abbott's formula. Different letters show the difference between isolates according to the LSD multiple comparison test ( $P < 0.05$ ). Bars show standard deviation. Ags1: *Bacillus cereus*, Ags2: *Bacillus* sp., Ags3: *Bacillus megaterium*, Ags4: *Enterobacter aerogenes*, Ags5: *Acinetobacter calcoaceticus*, Ags6: *Enterobacter* sp., Ags7: *Pseudomonas putida*, Ags8: *Enterococcus gallinarum*, Ags9: *Stenotrophomonas maltophilia*.

In all bioassays, the highest insecticidal infectivity determined in the *A. segetum* larvae was 60% for *Enterococcus gallinarum* (Ags8). The other isolates, Ags1, Ags2, Ags3, Ags4, Ags5, Ags6, Ags7, and Ags9, had 30%, 10%, 40%, 10%, 20%, 30%, 20%, and 30% mortality, respectively (Figure 2). The results of the bioassays indicated that all isolates (spore-forming and non-spore-forming) are pathogenic to the pest at different ratios, except for Ags2 and Ags4 (Figure 2). Lipa and Wiland (1972) observed the highest insecticidal activity on isolate *Pseudomonas fluorescens* 94-83a with 88% mortality by injection of bacterial suspension. The same strain showed 17%

mortality in *A. segetum* larvae by microfeeding. However, we determined the highest insecticidal activity in isolate Ags8 (*Enterococcus gallinarum*) 8 days after application by direct feeding of larvae with lettuce. That means that this isolate can be more pathogenic to *A. segetum* larvae under field conditions.

Enterococci are gram-positive and naturally occur in large numbers in the intestines of mammals, birds, reptiles, and insects (Lukasova and Sustackova 2003). These bacteria are not regarded as primary pathogens, but they are generally recognized as nosocomial pathogens worldwide (Linden and Miller 1999). Some studies have shown the isolation of *Enterococcus gallinarum* from several insect species (Xiang et al. 2006; Cox and Gilmore 2007). Although many Enterococci species are known as common symbionts in the gastrointestinal tracts of domestic animals, with this study, we showed for the first time that *Enterococcus gallinarum* (Ags8) has an important mortality value against *A. segetum*.

In conclusion, we determined members of the bacterial flora of *Agrotis segetum* larvae. Compared to previous studies of the bacterial flora of *A. segetum*, all bacterial isolates determined in this study were different. Isolate Ags8 appears to be a promising biocontrol agent for use against this pest. Further research will investigate the potential of these bacteria as a biocontrol agent against *A. segetum*.

### Acknowledgment

We would like to thank Dr. Christer Löfstedt and Erling Jirle for kindly providing *A. segetum* eggs (Department of Ecology, Lund University, Sweden).

### References

- Abbott WS (1925) A method of computing the effectiveness of an insecticide. *J Econ Entomol* 18: 265-267.
- Bahar AA, Demirbag Z (2007) Isolation of pathogenic bacteria from *Oberea linearis* (Coleoptera: Cerambycidae). *Biologia* 62: 13-18.
- Barry AL, Gavan TL, Badal RE, Telenson MJ (1982) Sensitivity, specificity, and reproducibility of the automicrobic system (with the *Enterobacteriaceae*-plus Biochemical Card) for identifying clinical isolates of Gram-negative bacilli. *J Clin Microbiol* 15: 582-588.
- Ben-Dov E, Boussiba S, Zaritsky A (1995) Mosquito larvicidal activity of *Escherichia coli* with combinations of genes from *Bacillus thuringiensis* subsp. *israelensis*. *J Bacteriol* 10: 2581-2587.
- Caballero P, Osuna-Vargas E, Santiago-Alvarez C (1991) Efficacy of a Spanish strain of *Agrotis segetum* granulosis virus (Baculoviridae) against *Agrotis segetum* Schiff. (Lepidoptera: Noctuidae) on corn. *J Appl Entomol* 112: 59-64.
- Claus M (1992) A standardized Gram staining procedure. *World J Microbiol Biotechnol* 8: 451-452.

- Cox CR, Gilmore MS (2007) Native microbial colonization of *Drosophila melanogaster* and its use as a model of *Enterococcus faecalis* pathogenesis. *Infect Immun* 75: 1565-1576.
- Demir I, Sezen K, Demirbag Z (2002) The first study on bacterial flora and biological control agent of *Anoplus roboris* (Suftr., Coleoptera). *J Microbiol* 40: 104-108.
- Hall TA (1999) Bioedit: A user-friendly biological sequence alignment editor and analysis program for windows 95/98/NT. *Nucleic Acids Symp* 41: 95-98.
- Ignoffo CM, Garcia C (1979) Susceptibility of larvae of the black cutworm to species of entomopathogenic bacteria, fungi, protozoa, and viruses. *J Econ Entomol* 72: 767-769.
- Ince IA, Kati H, Yilmaz H, Demir I, Demirbag Z (2008) Isolation and identification of bacteria from *Thaumetopoea pityocampa* Den. and Schiff. (Lepidoptera: Thaumetopoeidae) and determination of their biocontrol potential. *World J Microbiol Biotechnol* 24: 3005-3015.
- Jakubowska A, Van Oers MM, Ziemnicka J, Lipa JJ, Vlak JM (2005) Molecular characterization of *Agrotis segetum* nucleopolyhedrovirus from Poland. *J Invertebr Pathol* 90: 64-68.
- Kandler O, Weiss N (1986) Regular, nonsporng gram-positive rods. In: *Bergey's Manual of Systematic Bacteriology* (Eds. PHA Sneath, NS Mair, ME Sharpe, JG Holt), Williams and Wilkins, Baltimore, p. 1224.
- Kuzina LV, Peloquin JJ, Vacek DC, Miller TA (2001) Isolation and identification of bacteria associated with adult laboratory Mexican fruit flies, *Anastrepha ludens* (Diptera: Tephritidae). *Curr Microbiol* 42: 290-294.
- Lau WL, Jumars PA, Armbrust EV (2002) Genetic diversity of attached bacteria in the hindgut of the deposit-feeding shrimp *Neotrypaea* (Formerly *Callianassa*) Californiensis (Decapoda: Thalassinidae). *Microb Ecol* 43: 455-466.
- Linden PK, Miller CB (1999) Vancomycin-resistant *enterococci*: the clinical effect of a common nosocomial pathogen. *Diagn Microbiol Infect Dis* 33: 113-120.
- Lipa JJ, Wiland E (1972) Bacteria isolated from cutworms and their infectivity to *Agrotis* spp. (Lepitoptera: Noctuidae). *Acta Microbiol Pol* 4: 127-140.
- Lossbroek TG, Theunissen J (1985) The entomogenous nematode *Neoplectana bibionis* as a biological control agent of *Agrotis segetum* in lettuce. *Exp Appl Nematol* 39: 261-264.
- Lukasova J, Sustackova A (2003) *Enterococci* and antibiotic resistance. *Acta Vet Brno* 72: 315-323.
- The Ministry of Agriculture of Turkey (2008) *The Technical Guide of Agricultural Control*, Vol 3, Başak Publisher, Ankara, pp. 262-265.
- Mitchell FL, Smith JW (1985) Pathology and bioassays of the lesser cornstalk borer (*Elasmopalpus lignosellus*) entomopoxvirus. *J Invertebr Pathol* 45: 75-80.
- Murrell A, Dobson SJ, Yang X, Lacey E, Barker SC (2003) A survey of bacterial diversity in ticks, lice and fleas from Australia. *Parasitol Res* 89: 326-334.
- Ohba M, Aizawa K (1986) Distribution of *Bacillus thuringiensis* in soil of Japan. *J Invertebr Pathol* 47: 277-287.
- Osborn F, Berlioz L, Vitelli-Flores J, Monsalve W, Dorta B, Lemoine VR (2002) Pathogenic effects of bacteria isolated from larvae of *Hylesia metabus* Crammer (Lepidoptera: Saturniidae). *J Invertebr Pathol* 80: 7-12.
- Palleroni NJ (1984) Gram-negative aerobic rods and cocci, family I: *Pseudomonas*aceae. In: *Bergey's Manual of Systematic Bacteriology* (Eds. NR Krieg, JG Holt), Williams and Wilkins, Baltimore, pp. 141-199.
- Prescott LM, Harley JP, Klein DA (1996) *Microbiology*, Wm C Brown Publishers, England.
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, New York.
- Sandra WW, Douglas IG (2004) Microorganisms associated with field-collected *Chrysoperla rufilabris* (Neuroptera: Chrysopidae) adults with emphasis on yeast symbionts. *Biol Cont* 29: 155-168.
- Sezen K, Demir I, Demirbag Z (2005) Investigations on bacteria as a potential biological control agent of summer chafer, *Amphimallon solstitiale* L. (Coleoptera: Scarabaeidae). *J Microbiol* 43: 463-468.
- Soutourina OA, Semenova EA, Parfenova VV, Danchin A, Bertin P (2001) Control of bacterial motility by environmental factors in polarly elagellated and peritrichous bacteria isolated from Lake Baikal. *Appl Environ Microbiol* 67: 3852-3859.
- Steenberg T, Ogaard L (2000) Mortality in hibernating turnip moth larvae, *Agrotis segetum*, caused by *Tolyposcladium cylindrosporium*. *Mycol Res* 104: 87-91.
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 24: 1596-1599.
- Thiery I, Frachon E (1997) Identification, isolation, culture and preservation of entomopathogenic bacteria. In: *Manual of Techniques in Insect Pathology* (Ed. AL Lacey), Academic Press, London, pp. 55-73.
- Turchaninova LP (1972) The bacterial flora of healthy *Agrotis segetum* Schiff in various developmental stages. *Mikrobiol Zh* 34: 119-20.
- Turchaninova LP (1973) Pathogenic bacterial flora of the winter cutworm (*Agrotis segetum* Schiff). *Mikrobiol Zh* 35: 576-80.
- William GW, Susan MB, Dale AP, David JL (1991) 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol* 173: 697-703.
- Xiang H, Wei GF, Jia S, Huang J, Miao XX, Zhou Z, Zhao LP, Huang YP (2006) Microbial communities in the larval midgut of laboratory and field populations cotton bollworm (*Helicoverpa armigera*). *Can J Microbiol* 52: 1085-1092.
- Yilmaz H, Sezen K, Kati H, Demirbag Z (2006) The first study on the bacterial flora of the European spruce bark beetle, *Dendroctonus micans* (Coleoptera: Scolytidae). *Biologia* 61: 679-686.
- Yun JS, Kim HH, Kim DW, Lee SM, Kim DS, Lee DW (2004) Pathogenicities of entomopathogenic fungi, *Beauveria bassiana* and *Metarhizium anisopliae* against lepidopterous insect pests, *Agrotis segetum*, *Artogeia rapae*, *Mamestra brassicae*, *Plutella xylostella*, *Spodoptera exigua* and *Spodoptera litura*. *Kor Turfgrass Sci* 18: 221-229.
- Zethner O (1980) Control of *Agrotis segetum* (Lepidoptera: Noctuidae) root crops by granulosis virus. *Entomophaga* 25: 27-35.